

On Ergodicity, Infinite Flow and Consensus in Random Models

Behrouz Touri and Angelia Nedić

Abstract

We consider the ergodicity and consensus problem for a discrete-time linear dynamic model driven by random stochastic matrices, which is equivalent to studying these concepts for the product of such matrices. Our focus is on the model where the random matrices have independent but time-variant distribution. We introduce a new phenomenon, the infinite flow, and we study its fundamental properties and relations with the ergodicity and consensus. The central result is the infinite flow theorem establishing the equivalence between the infinite flow and the ergodicity for a class of independent random models, where the matrices in the model have a common steady state in expectation and a feedback property. For such models, this result demonstrates that the expected infinite flow is both necessary and sufficient for the ergodicity. The result is providing a deterministic characterization of the ergodicity, which can be used for studying the consensus and average consensus over random graphs.

Index Terms

Ergodicity, random consensus, linear random model, product of random matrices.

I. INTRODUCTION

There is an evidence of growing number of applications in decentralized control of networked agents, as well as social and other networks where the consensus is used as a mechanism for decentralized coordination of agent actions. The focus of this paper is on a canonical consensus problem for a linear discrete-time dynamic system driven by a general model of random matrices, where the matrices are row-stochastic. Currently, investigating whether the model reaches a

Department of Industrial and Enterprise Systems Engineering, University of Illinois, Urbana, IL 61801, Email: {touri1,angelia}@illinois.edu. This research is supported by the National Science Foundation under CAREER grant CMMI 07-42538.

consensus or not is often done by exploring the conditions that ensure the ergodicity of the model, which in turn always guarantees the consensus.

In this paper, we propose an alternative approach by introducing a concept of the model with infinite flow, which can be interpreted as infinite information flow over time between a group of agents and the other agents in the network. We show that the infinite flow property is closely related to the ergodicity and, hence, to the consensus. In particular, we show the equivalence between the infinite flow and the ergodicity for a class of independent random models.

We start by a comprehensive study of the fundamental relations and properties of the ergodicity, consensus and infinite flow for general random models, independent models and independent identically distributed (i.i.d.) random models. We then investigate the random models with a feedback property and the models with a common steady state for the expected matrices. Both of these properties have been used in the analysis of consensus models, but a deeper understanding of their roles has not been observed. We classify feedback property in three basic types from weak to strong and show some relations for them. We then study the models that have a common steady state in expectation. By combining all the pieces together, we show that the ergodicity of the model is equivalent to the infinite flow property for a class of independent random models with feedback property and a common steady state in expectation, as given in infinite flow theorem (Theorem 7). The infinite flow theorem also establishes the equivalence between the infinite flow property of the model and the infinite flow property of the expected model. Furthermore, the theorem also establishes the equivalence between the ergodicity of the model and the ergodicity of the expected model. As such, this theorem provides a novel deterministic characterization of the ergodicity, thus rendering another tool for studying the consensus over random networks and convergence of random consensus algorithms.

The main contributions of this paper include: 1) the equivalence of the ergodicity of the model and the expected model for a class of independent random models with a feedback property and a common steady state in expectation; 2) the new insights and understanding of the ergodicity and consensus events over random networks brought to light through a new phenomena of infinite flow event, which to the best of our knowledge has not been known prior to this work; 3) novel comprehensive study of the fundamental properties of the consensus and ergodicity events for general class of independent random models; 4) new insights into the role of feedback property and the role of a common steady state in expectation for the ergodicity and consensus.

The study of the random product of stochastic matrices dates back to the early work in [32] where the convergence of the product of i.i.d. random stochastic matrices was studied using the algebraic and topological structures of the set of stochastic matrices. This work was further extended in [22], [23], [10] by using results from ergodic theory of stationary processes and their algebraic properties. In [33], the ergodicity and consensus of the product of i.i.d. random stochastic matrices was studied using tools from linear algebra and probability theory, and a necessary and sufficient condition for the ergodicity was established for a class of i.i.d. models. Independently, the same problem was tackled in [13], where an exponential convergence bound was established. Recently, the work in [33] was extended to ergodic stationary processes in [34].

In all of the works [32], [22], [23], [10], [33], [13], [34], the underlying random models are assumed to be either i.i.d. or stationary processes, both of which imply time-invariant distribution on the random model. Unlike these works, our work in this paper is focused on the independent random models with *time-variant distributions*. Furthermore, we study ergodicity and consensus for such models using martingale and supermartingale convergence results combined with the basic tools from probability theory. Our work is also related to the consensus over random networks [14], optimization over random networks [21], and the consensus over a network with random link failures [29]. Related are also gossip and broadcast-gossip schemes giving rise to a random consensus over a given connected bi-directional communication network [6], [11], [1], [30]. On a broader basis, the paper is related to the literature on the consensus over networks with noisy links [15], [16], [18], [35] and the deterministic consensus in decentralized systems models [36], [37], [38], [2], [20], [17], [27], [28], [31] including the effects of quantization and delay [19], [24], [8], [7], [4], [30].

The paper is organized as follows. In Section II, we describe a discrete-time random linear dynamic system of our interest, and introduce the ergodicity, consensus and infinite flow events. In Section III, we explore the relations among these events and establish their 0-1 law and other properties by considering general, independent and i.i.d. random models. In Section IV, we discuss models with feedback properties and provide classification of such properties with insights into their relations. We also consider independent random model with a common steady state in expectation. In Section V, we focus on independent random models with infinite flow property. We establish necessary and sufficient conditions for ergodicity, and briefly discuss some implications of these conditions. We conclude in Section VI.

Notation and Basic Terminology. We view all vectors as columns. For a vector x , we write x_i to denote its i th entry, and we write $x \geq 0$ ($x > 0$) to denote that all its entries are nonnegative (positive). We use x^T to denote the transpose of a vector x . We write $\|x\|$ to denote the standard Euclidean vector norm i.e., $\|x\| = \sqrt{\sum_i x_i^2}$. We use e_i to denote the vector with the i th entry equal to 1 and all other entries equal to 0. We write e to denote the vector with all entries equal to 1. A vector a is *stochastic* when $a \geq 0$ and $\sum_i a_i = 1$. We write $\{x(k)\}$ or $\{x(k)\}_{k \geq 0}$ to denote a sequence $x(0), x(1), \dots$ of some elements, and we write $\{x(k)\}_{k \geq t}$ to denote the truncated sequence $x(t), x(t+1), \dots$ for $t > 0$. For a given set C and a subset S of C , we write $S \subset C$ to denote that S is a proper subset of C . A set $S \subset C$ such that $S \neq \emptyset$ is referred to as a *nontrivial* subset of C . We write $[m]$ to denote the integer set $\{1, \dots, m\}$. For a set $S \subset [m]$, we let \bar{S} denote the complement set of S with respect to $[m]$, i.e., $\bar{S} = \{i \in [m] \mid i \notin S\}$.

We denote the identity matrix by I . For a vector v , we use $\text{diag}(v)$ to denote the diagonal matrix with diagonal entries being the components v_i of the vector v . For a matrix W , we write W_{ij} to denote its (i, j) th entry, W^i to denote its i th column vector, and W^T to denote its transpose. For an $m \times m$ matrix W , we use $\sum_{i < j} W_{ij}$ to denote the summation of the entries W_{ij} over all $i, j \in [m]$ with $i < j$. A matrix W is *row-stochastic* when its entries are nonnegative and the sum of the row entries is 1 for each row. Since we deal exclusively with row-stochastic matrices, we will refer to such matrices simply as *stochastic*. We let \mathbb{S}^m denote the set of $m \times m$ stochastic matrices. A matrix W is *doubly stochastic* when both W and W^T are stochastic.

We write $E[X]$ and $\text{Var}(X)$ to denote, respectively, the expected value and the variance of a random variable X . For an event \mathcal{A} , we use $\Pr(\mathcal{A})$ to denote its probability and $1_{\mathcal{A}}$ to denote its characteristic function. If $\Pr(\mathcal{A}) = 1$, we say that the event \mathcal{A} happens almost surely. We often abbreviate “almost surely” by *a.s.*

II. PROBLEM FORMULATION AND TERMINOLOGY

Throughout this article, we deal exclusively with the matrices in the set \mathbb{S}^m , the set of $m \times m$ stochastic matrices. We consider the topology induced by the open sets in \mathbb{S}^m with respect to the Euclidean norm and the Borel sigma-algebra $\mathcal{F}_{\mathbb{S}^m}$ of this topology. We assume that we are given a probability space $(\Omega, \mathcal{R}, \Pr(\cdot))$ and a measurable function $W : \Omega \rightarrow \prod_{k=0}^{\infty} \mathbb{S}^m$. To every $\omega \in \Omega$, the function $W(\cdot)$ is assigning a discrete time process $\{W(k)\}(\omega)$ in the countable product measurable space $\prod_{k=0}^{\infty} (\mathbb{S}^m, \mathcal{F}_{\mathbb{S}^m})$, where $W(k) \in \mathbb{S}^m$ is the random matrix of the

process at time k . We refer to the process $\{W(k)\}(\omega)$ interchangeably as a *random chain* or a *random model* and, when suitable, we suppress the explicit dependence on the variable ω . We say that the chain $\{W(k)\}$ is independent if the sigma algebra generated by the $W(k)$ s for different $k \geq 0$ are independent. Furthermore, if $W(k)$ s are identically distributed, we say that the model is independently identically distributed (i.i.d.).

With a given random chain $\{W(k)\}$, we associate a linear discrete-time dynamic system of the following form:

$$x(k+1) = W(k)x(k) \quad \text{for } k = 0, 1, 2, \dots, \quad (1)$$

where $x(k) \in \mathbb{R}^m$ is a state vector at time k and $x(0)$ is the initial state vector. We will often refer to the system in (1) as *the dynamic system driven by the chain $\{W(k)\}$* .

We are interested in providing conditions guaranteeing that the dynamic system reaches a consensus almost surely. Since reaching the consensus is closely related to the ergodicity of the chain, we are also interested in studying the ergodicity on the fundamental level. In our study of the random consensus and ergodicity, we use another property of the chain, an infinite flow property. We start by providing these basic notions for a deterministic chain.

Definition 1: Given a deterministic chain $\{G(k)\} \subset \mathbb{S}^m$, we say that:

The system $z(k+1) = G(k)z(k)$ *reaches a consensus* if for any initial state $z(0) \in \mathbb{R}^m$, there exists a scalar $c(z(0))$ such that $\lim_{k \rightarrow \infty} \|z(k) - c(z(0))e\| = 0$.

The chain $\{G(k)\}$ is *ergodic* if for any $t \geq 0$ and $j \in [m]$, there is a scalar $g_j(t)$ such that

$$\lim_{k \rightarrow \infty} \Phi_{ij}(k, t) = g_j(t) \quad \text{for all } i \in [m],$$

where $\Phi(k, t) = G(k)G(k-1) \cdots G(t)$ for $k > t$ and $t \geq 0$.

The chain $\{G(k)\}$ *has infinite flow property* if $\sum_{k=0}^{\infty} \sum_{i \in S, j \in \bar{S}} (G_{ij}(k) + G_{ji}(k)) = \infty$ for any nontrivial subset $S \subset [m]$.

In the definition of the infinite flow property, the quantity $\sum_{i \in S, j \in \bar{S}} (G_{ij}(k) + G_{ji}(k))$ can be interpreted as a flow between the subset S and its complement \bar{S} in a weighted graph. In particular, consider the undirected weighted graph $\mathcal{G}(k)$ with the node set $[m]$, the edge set induced by the positive entries in $G(k) + G^T(k)$, and the weight matrix $G(k) + G^T(k)$. In this case, the quantity $\sum_{i \in S, j \in \bar{S}} (G_{ij}(k) + G_{ji}(k))$ represents the flow in graph $\mathcal{G}(k)$ across the cut (S, \bar{S}) for a nontrivial node set $S \subset [m]$ and its complement \bar{S} . For the graphs $\mathcal{G}(k)$ induced

by the matrices $G(k)$, the infinite flow property requires that the total flow in time across any nontrivial cut (S, \bar{S}) is infinite, which could be viewed as infinite information exchange between the nodes in S and \bar{S} .

The ergodicity is equivalent to the following condition [9]: for any $t \geq 0$ and $x \in \mathbb{R}^m$, there is a scalar $\gamma(t, x)$ such that $\lim_{k \rightarrow \infty} \Phi(k, t)x = \gamma(t, x)e$. Since the matrices $G(k)$ have finite dimension, the ergodicity is also equivalent to the following condition: for any $t \geq 0$ and any $\ell \in [m]$, there is a scalar $\gamma_\ell(t)$ such that $\lim_{k \rightarrow \infty} \Phi(k, t)e_\ell = \gamma_\ell(t)e$. Similarly, due to the linearity and finite dimension of the system $z(k+1) = G(k)z(k)$, the consensus can be studied by considering only the initial states $x(0) = e_\ell$ with $\ell \in [m]$ rather than all $x(0) \in \mathbb{R}^m$.

Clearly, the ergodicity of the chain implies reaching a consensus. However, a consensus may be reached even if the chain $\{G(k)\}$ is not ergodic, as seen in the following example.

Example 1: Let $G(0) = ev^T$ for a stochastic vector v , and let $G(k) = I$ for all $k \geq 1$. Then, we have $\Phi(k, 0) = ev^T$ for all $k \geq 1$, implying that the system $x(k+1) = G(k)x(k)$ reaches a consensus. However, the chain $\{G(k)\}$ is not ergodic since $\Phi(k, t) = I$ for any $k > t \geq 1$. \square

Using Definition 1, we now introduce the corresponding events of consensus, ergodicity and infinite flow. Given a random chain $\{W(k)\}$, let \mathcal{C} denote the event that the system in (1) reaches a consensus for any initial state $x(0)$. Let \mathcal{E} denote the event that the chain $\{W(k)\}$ is ergodic, and let \mathcal{F} denote the event that the chain has the *infinite flow property*. We refer to \mathcal{C} , \mathcal{E} and \mathcal{F} as the *consensus event*, the *ergodicity event* and the *infinite flow event*, respectively. We say that the *model is ergodic* if the ergodicity event \mathcal{E} occurs almost surely. The *model admits consensus* if the consensus event \mathcal{C} occurs almost surely. The *model has infinite flow* if the infinite flow event \mathcal{F} occurs almost surely. The *model has expected infinite flow* if its expected chain $\{E[W(k)]\}$ has infinite flow.

III. INFINITE FLOW, ERGODICITY AND CONSENSUS

In this section, we further study the ergodicity event \mathcal{E} , the consensus event \mathcal{C} and the infinite flow event \mathcal{F} under different assumptions on the nature of the randomness in the model. In particular, in Section III-A we establish some fundamental relations among \mathcal{E} , \mathcal{C} and \mathcal{F} . In Section III-B, we investigate the 0-1 law properties of these events, while in Section III-C we provide some relations for a random model and its expected model.

Throughout the rest of the paper, we use the following notation. For a stochastic matrix W and a nontrivial subset $S \subset [m]$, we define W_S as follows:

$$W_S = \sum_{i \in S, j \in \bar{S}} (W_{ij} + W_{ji}). \quad (2)$$

Note that for a given a random model $\{W(k)\}$, the infinite flow event \mathcal{F} is given by

$$\mathcal{F} = \bigcap_{S \subset [m]} \left\{ \sum_{k=0}^{\infty} W_S(k) = \infty \right\}. \quad (3)$$

Note that the infinite flow event requires that, across any nontrivial cut (S, \bar{S}) , the total flow in the random (undirected) graphs induced by the matrices $W(k) + W^T(k)$, $k \geq 0$, is infinite.

A. Basic Relations

As discussed in Section II, we have $\mathcal{E} \subseteq \mathcal{C}$ for any random model. We here show that the ergodicity event is, also, always contained in the infinite flow event, i.e., $\mathcal{E} \subseteq \mathcal{F}$. We establish this by using the following result applicable to a deterministic dynamic system.

Lemma 1: Let $\{A(k)\} \subset \mathbb{S}^m$ be a deterministic sequence, and let $\{z(k)\}$ be generated by $z(k+1) = A(k)z(k)$ for all $k \geq 0$ with an initial state $z(0) \in \mathbb{R}^m$. Then, for any nontrivial subset $S \subset [m]$ and $k \geq 0$, we have

$$\begin{aligned} \max_{i \in S} z_i(k+1) &\leq \max_{s \in S} z_s(0) + d(z(0)) \sum_{t=0}^k A_S(t), \\ \min_{j \in \bar{S}} z_j(k+1) &\geq \min_{r \in \bar{S}} z_r(0) - d(z(0)) \sum_{t=0}^k A_S(t), \end{aligned}$$

where $d(y) = \max_{\ell \in [m]} y_\ell - \min_{r \in [m]} y_r$ for $y \in \mathbb{R}^m$.

Proof: Let $S \subset [m]$ be an arbitrary nontrivial set and let $k \geq 0$ be arbitrary. Let $z_{\min}(k) = \min_{r \in [m]} z_r(k)$ and $z_{\max}(k) = \max_{s \in [m]} z_s(k)$. Since $z_i(k+1) = \sum_{\ell=1}^m A_{i\ell}(k) z_\ell(k)$, by the stochasticity of $A(k)$ we have $z_i(k) \in [z_{\min}(0), z_{\max}(0)]$ for all $i \in [m]$ and all k . Then, we obtain for $i \in S$,

$$z_i(k+1) = \sum_{\ell \in S} A_{i\ell}(k) z_\ell(k) + \sum_{\ell \in \bar{S}} A_{i\ell}(k) z_\ell(k) \leq \sum_{\ell \in S} A_{i\ell}(k) \left(\max_{s \in S} z_s(k) \right) + z_{\max}(0) \sum_{\ell \in \bar{S}} A_{i\ell}(k),$$

where the inequality follows by $A_{i\ell}(k) \geq 0$. By the stochasticity of $A(k)$, we also obtain

$$z_i(k+1) \leq \left(1 - \sum_{\ell \in \bar{S}} A_{i\ell}(k) \right) \max_{s \in S} z_s(k) + z_{\max}(0) \sum_{\ell \in \bar{S}} A_{i\ell}(k)$$

$$= \max_{s \in S} z_s(k) + \left(z_{\max}(0) - \max_{s \in S} z_s(k) \right) \sum_{\ell \in \bar{S}} A_{i\ell}(k).$$

By the definition of A_S in (2), we have $0 \leq \sum_{\ell \in \bar{S}} A_{i\ell}(k) \leq A_S(k)$. Since $z_{\max}(0) - \max_{s \in S} z_s(k) \geq 0$, it follows

$$z_i(k+1) \leq \max_{s \in S} z_s(k) + (z_{\max}(0) - \max_{s \in S} z_s(k)) A_S(k) \leq \max_{s \in S} z_s(k) + d(z(0)) A_S(k),$$

where the last inequality holds since $z_{\max}(0) - \max_{s \in S} z_s(k) \leq z_{\max}(0) - z_{\min}(0) = d(z(0))$.

By taking the maximum over all $i \in S$ in the preceding relation and by recursively using the resulting inequality, we obtain

$$\max_{i \in S} z_i(k+1) \leq \max_{s \in S} z_s(k) + d(z(0)) A_S(k) \leq \cdots \leq \max_{s \in S} z_s(0) + d(z(0)) \sum_{t=0}^k A_S(t).$$

The relation for $\min_{j \in \bar{S}} z_j(k+1)$ follows from the preceding relation by considering $\{z(k)\}$ generated with the starting point $-z(0)$. ■

Using Lemma 1, we now show that the ergodicity event is contained in the infinite flow event.

Theorem 1: Let $\{A(k)\} \subset \mathbb{S}^m$ be an ergodic deterministic chain. Then $\sum_{k=0}^{\infty} A_S(k) = \infty$ for any nontrivial $S \subset [m]$. In particular, we have $\mathcal{E} \subseteq \mathcal{F}$ for any random model.

Proof: To arrive at a contradiction, assume that there is a nontrivial set $S \subset [m]$ such that $\sum_{k=0}^{\infty} A_S(k) < \infty$. Since the matrices $A(k)$ are stochastic, we have $A_S(k) \geq 0$ for all k . Therefore, there exists large enough $\bar{t} \geq 0$ such that

$$\sum_{k=\bar{t}}^{\infty} A_S(k) < \frac{1}{4}. \quad (4)$$

Now, define the vector $\bar{z} = (\bar{z}_1, \dots, \bar{z}_m)^T$ as follows:

$$\bar{z}_i = \begin{cases} 0 & \text{if } i \in S, \\ 1 & \text{if } i \in \bar{S}, \end{cases}$$

and consider the dynamic system

$$z(k+1) = A(k)z(k) \quad \text{for } k \geq \bar{t},$$

which is started at time \bar{t} in state $z(\bar{t}) = \bar{z}$. Note that Lemma 1 applies to the case where the time $t = \bar{t}$ is taken as initial time, in which case $d(0)$ corresponds to $d(\bar{t}) = \max_i z_i(\bar{t}) - \min_j z_j(\bar{t})$.

Also, note that $d(\bar{t}) = 1$ by the definition of the starting state \bar{z} . Thus, by applying Lemma 1, we have for all $k \geq \bar{t}$,

$$\begin{aligned}\max_{i \in S} z_i(k+1) &\leq \max_{s \in S} z_s(\bar{t}) + \sum_{t=\bar{t}}^k A_S(t), \\ \min_{j \in \bar{S}} z_j(k+1) &\geq \min_{r \in \bar{S}} z_r(\bar{t}) - \sum_{t=\bar{t}}^k A_S(t).\end{aligned}$$

Since $\max_{s \in S} z_s(\bar{t}) = 0$ and $\min_{r \in \bar{S}} z_r(\bar{t}) = 1$, it follows

$$\max_{i \in S} z_i(k+1) \leq \sum_{t=\bar{t}}^k A_S(t), \quad \min_{j \in \bar{S}} z_j(k+1) \geq 1 - \sum_{t=\bar{t}}^k A_S(t).$$

From the preceding relations and Eq. (4) we have

$$\liminf_{k \rightarrow \infty} (z_j(k) - z_i(k)) > 1 - 2 \sum_{t=\bar{t}}^{\infty} A_S(t) = \frac{1}{2} \quad \text{for any } j \in \bar{S} \text{ and } i \in S,$$

thus showing that the chain $\{A(k)\}$ is not ergodic - a contradiction. Therefore, we must have $\sum_{t=0}^{\infty} A_S(t) = \infty$ for any nontrivial $S \subset [m]$.

Based on the preceding argument, and the definitions of \mathcal{E} and \mathcal{F} , for any random model $\{W(k)\}$ we conclude that $\omega \in \mathcal{E}$ implies $\omega \in \mathcal{F}$. Hence, $\mathcal{E} \subseteq \mathcal{F}$ for any random model. ■

Theorem 1 shows that an ergodic model must have an infinite flow property. In other words, the infinite flow property of any random model is necessary for the ergodicity of the model. Later in Theorem 6, for a certain class of random models, we will show that the infinite flow is also sufficient for the ergodicity.

Figure 1 illustrates the inclusions $\mathcal{E} \subseteq \mathcal{C}$ and $\mathcal{E} \subseteq \mathcal{F}$ for a general random model. The inclusion $\mathcal{E} \subseteq \mathcal{C} \cap \mathcal{F}$ in Figure 1 can be strict as seen in the following example.

Example 2: Consider the 2×2 chain $\{A(k)\}$ defined by

$$A(0) = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix} \quad \text{and} \quad A(k) = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \text{for } k \geq 1.$$

For any stochastic matrix B , we have $BA(0) = A(0)$ and hence, the model admits consensus. Furthermore, the model has infinite flow property. However, the chain $\{A(k)\}_{k \geq 1}$ does not admit consensus. Therefore, in this case $\mathcal{C} \cap \mathcal{F} = \Omega$ (the entire space of realizations) while $\mathcal{E} = \emptyset$. □

Next, we provide a sufficient condition for the ergodicity and consensus events to coincide.

Lemma 2: Let $\{W(k)\}$ be a (not necessarily independent) random chain such that each matrix $W(k)$ is invertible almost surely. Then, we have $\mathcal{E} = \mathcal{C}$ almost surely.




Fig. 1. The relations among the ergodicity, consensus, and infinite flow events for a general random model.

Proof: The inclusion $\mathcal{E} \subseteq \mathcal{C}$ follows from the definition, so it suffices to show $\mathcal{C} \subseteq \mathcal{E}$ almost surely. In turn, to show $\mathcal{C} \subseteq \mathcal{E}$ almost surely, it suffices to prove that $\mathcal{C} \cap R \subseteq \mathcal{E}$ for a set R such that $\Pr(R) = 1$. For each $k \geq 0$, let R_k be the set of instances ω such that the matrix $W(k)(\omega)$ is invertible. Define $R = \bigcap_{k=0}^{\infty} R_k$. We have $\Pr(R_k) = 1$ for each $k \geq 0$ by our assumption that each matrix $W(k)$ is invertible almost surely. In view of this and the fact that the collection $\{R_k\}$ is countable, it follows that $\Pr(R) = 1$.

We next show that $\mathcal{C} \cap R \subseteq \mathcal{E}$. Let $\omega \in \mathcal{C} \cap R$ so that $W(k)(\omega)$ has full rank for all $k \geq 0$. To simplify notation, let $\tilde{W}(k) = W(k)(\omega)$. Consider an arbitrary starting time $s \geq 0$. We show that the consensus is reached for the dynamic $z(t) = \tilde{W}(t-1)z(t-1)$ with $t > s$, i.e., for any $z(s) \in \mathbb{R}^m$, we have $\lim_{t \rightarrow \infty} z(t) = ce$ for some $c \in \mathbb{R}$. For a given $z(s) \in \mathbb{R}^m$, define $\tilde{x}(0) = [\tilde{W}(s-1) \cdots \tilde{W}(1)\tilde{W}(0)]^{-1}z(s)$ and consider the dynamic $\tilde{x}(k) = \tilde{W}(k-1)\tilde{x}(k-1)$ started at time $t_0 = 0$ with the initial vector $\tilde{x}(0)$. By the definition of $\tilde{x}(0)$, we have $\tilde{x}(s) = \tilde{W}(s-1) \cdots \tilde{W}(1)\tilde{W}(0)\tilde{x}(0) = z(s)$. Therefore, for all $t > s$,

$$\tilde{x}(t) = \tilde{W}(t-1) \cdots \tilde{W}(s)\tilde{W}(s-1) \cdots \tilde{W}(1)\tilde{W}(0)\tilde{x}(0) = \tilde{W}(t-1) \cdots \tilde{W}(s)z(s) = z(t).$$

Since $\omega \in \mathcal{C}$, by the definition of $\tilde{W}(k)$ we have $\lim_{t \rightarrow \infty} \tilde{x}(t) = ce$ for some $c \in \mathbb{R}$. In view of the preceding relation, it follows that $\lim_{t \rightarrow \infty} z(t) = ce$, thus showing that the dynamic system $z(t) = \tilde{W}(t-1)z(t-1)$, $t > s$, reaches a consensus. Since this is true for arbitrary $s \geq 0$ and $z(s) \in \mathbb{R}^m$, the chain $\{W(k)\}(\omega)$ is ergodic, which implies $\omega \in \mathcal{E}$. ■

In general, there may be no further refinements of inclusion relations among the events \mathcal{E} , \mathcal{F} and \mathcal{C} even when the model is independent, as indicated by the following example.

Example 3: Consider an independent random model where for $p \in (0, 1]$, we have

$$W(0) = \frac{1}{m} ee^T \text{ with probability } p \quad \text{and} \quad W(0) = I \text{ with probability } 1 - p,$$

and $W(k) = I$ with probability 1 for all $k \geq 1$. In this case, the consensus event \mathcal{C} happens with probability $p > 0$. However, the infinite flow and the ergodicity events are empty sets. □

Example 3 shows that we can have $\mathcal{C} \cap \mathcal{E} = \emptyset$ and $\mathcal{C} \cap \mathcal{F} = \emptyset$, while $\mathcal{C} \neq \emptyset$. Thus, even for an independent model the consensus event need not be contained in either \mathcal{E} or \mathcal{F} . However, if we further restrict our attention to i.i.d. models, we can show that $\mathcal{E} = \mathcal{C}$ almost surely. To establish this, we make use of the following lemma.

Lemma 3: Let $A \in \mathbb{S}^m$ and $x \in \mathbb{R}^m$, and let $[Ax]_\ell$ denote the ℓ th component of the vector Ax . Assume that A is such that $\max_{i \in [m]} [Ae_\ell]_i - \min_{j \in [m]} [Ae_\ell]_j \leq \frac{1}{2m}$ for any $\ell \in [m]$. Then, we have $\max_i [Ax]_i - \min_j [Ax]_j \leq \frac{1}{2}$ for any $x \in [0, 1]^m$.

Proof: Let $x \in \mathbb{R}^m$ with $x_\ell \in [0, 1]$ for any $\ell \in [m]$. Then, we have for any $i, j \in [m]$,

$$y_i - y_j = \sum_{\ell=1}^m (A_{i\ell} - A_{j\ell})x_\ell \leq \sum_{\ell=1}^m |A_{i\ell} - A_{j\ell}| = \sum_{\ell=1}^m |[Ae_\ell]_i - [Ae_\ell]_j|.$$

By the assumption on A , we obtain $|[Ae_\ell]_i - [Ae_\ell]_j| \leq \max_{i \in [m]} [Ae_\ell]_i - \min_{j \in [m]} [Ae_\ell]_j \leq \frac{1}{2m}$. Hence, $y_i - y_j \leq \sum_{\ell=1}^m \frac{1}{2m} = \frac{1}{2}$, implying $\max_i y_i - \min_j y_j \leq \frac{1}{2}$. ■

We now provide our main result for i.i.d. models, which states that the ergodicity and the consensus events are almost surely equal. We establish this result by using Lemma 3 and the Borel-Cantelli lemma (see [12], page 50).

Theorem 2: We have $\mathcal{E} = \mathcal{C}$ almost surely for any i.i.d. random model.

Proof: Since $\mathcal{E} \subseteq \mathcal{C}$, the assertion is true when consensus occurs with probability 0. Therefore, it suffices to show that if the consensus occurs with a probability p other than 0, the two events are almost surely equal. Let $\Pr(\mathcal{C}) = p$ with $p \in (0, 1]$. Then, for all $\omega \in \mathcal{C}$,

$$\lim_{k \rightarrow \infty} d(x(k))(\omega) = 0 \quad \text{where} \quad d(x) = \max_i x_i - \min_j x_j,$$

and $\{x(k)\}(\omega)$ is the sequence generated by the dynamic system (1) with any $x(0) \in \mathbb{R}^m$.

For each $\ell \in [m]$, let $\{x^\ell(k)\}$ be the random sequence that is generated by the dynamic system in (1) with $x(0) = e_\ell$. Then, for any $\omega \in \mathcal{C}$, there is the smallest integer $K^\ell(\omega) \geq 0$ such that

$$d(x^\ell(k))(\omega) \leq \frac{1}{2m} \quad \text{for all } k \geq K^\ell(\omega).$$

Note that $d(x^\ell(k))(\omega)$ is a nonincreasing sequence (of k) for each $\ell \in [m]$. Hence, by letting $K(\omega) = \max_{\ell \in [m]} K^\ell(\omega)$ we obtain $d(x^\ell(k))(\omega) \leq \frac{1}{2m}$ for all $\ell \in [m]$ and $k \geq K(\omega)$. Thus, by applying Lemma 3, we have for almost all $\omega \in \mathcal{C}$,

$$d(x(k))(\omega) \leq \frac{1}{2} \quad \text{for all } k \geq K(\omega) \text{ and } x(0) \in [0, 1]^m. \quad (5)$$

By the definition of consensus, we have $\lim_{N \rightarrow \infty} \Pr(K \leq N) \geq \Pr(\mathcal{C}) = p$. Thus, by the continuity of the measure, there exists an integer N_1 such that $\Pr(K < N_1) \geq \frac{p}{2}$.

Now, let time $T \geq 0$ be arbitrary, and let l_k^T denote the N_1 -tuple of the matrices $W(s)$ driving the system (1) for $s = T + N_1k, \dots, T + N_1(k+1) - 1$ and $k \geq 0$, i.e.,

$$l_k^T = (W(T + N_1k), W(T + N_1k + 1), \dots, W(T + N_1(k+1) - 1)) \quad \text{for all } k \geq 0.$$

Let L_N denote the collection of all N -tuples (A_1, \dots, A_N) of matrices $A_i \in \mathbb{S}^m$, $i = 1, \dots, N$ such that for $x(N) = A_N A_{N-1} \dots A_1 x(0)$ with $x(0) \in [0, 1]^m$, we have $d(x(N)) \leq \frac{1}{2}$. By the definitions of l_k^T and L_N , relation (5) and relation $\Pr(K < N_1) \geq \frac{p}{2}$ state that $\Pr(\{l_0^T \in L_{N_1}\}) \geq \frac{p}{2}$. By the i.i.d. property of the model, the events $\{l_k^T \in L_{N_1}\}$, $k \geq 0$, are i.i.d. and the probability of their occurrence is equal to $\Pr(\{l_0^T \in L_{N_1}\})$, implying that $\Pr(\{l_k^T \in L_{N_1}\}) \geq \frac{p}{2}$ for all $k \geq 0$. Consequently, $\sum_{k=0}^{\infty} \Pr(\{l_k^T \in L_{N_1}\}) = \infty$. Since the events $\{l_k^T \in L_{N_1}\}$ are i.i.d., by Borel-Cantelli lemma $\Pr(\{\omega \in \Omega \mid \omega \in \{l_k^T \in L_{N_1}\} \text{ i.o.}\}) = 1$, where *i.o.* stands for infinitely often. Observing that the event $\{\omega \in \Omega \mid \omega \in \{l_k^T \in L_{N_1}\} \text{ i.o.}\}$ is contained in the consensus event for the chain $\{W(T+k)\}_{k \geq 0}$, we see that the consensus event for the chain $\{W(T+k)\}_{k \geq 0}$ occurs almost surely. Since this is true for arbitrary $T \geq 0$ it follows that the chain $\{W(k)\}$ is ergodic *a.s.*, implying $\mathcal{C} \subseteq \mathcal{E}$ *a.s.* This and the inclusion $\mathcal{E} \subseteq \mathcal{C}$ yield $\mathcal{C} = \mathcal{E}$ *a.s.* ■

Theorem 2 extends the equivalence result between the consensus and ergodicity for i.i.d. models given in Theorem 3.a and Theorem 3.b of [33] (and hence Corollary 4 in [33]), which are established there assuming that the matrices have positive diagonal entries almost surely. The relations among \mathcal{C} , \mathcal{E} , and \mathcal{F} for i.i.d. case are illustrated in Figure 2.

B. 0-1 Laws

In this section, we discuss 0-1 laws for the events \mathcal{E} , \mathcal{F} and \mathcal{C} for independent random models. The 0-1 laws specify the trivial events, which are the events occurring with either probability 0 or 1. The ergodicity event is a trivial event, as shown¹ in [33], Lemma 1. Since the ergodicity event is always contained in the consensus event, the ergodicity event \mathcal{E} occurs with probability 0 whenever the consensus event \mathcal{C} occurs with a probability $p \in (0, 1)$. In other words, we may have $\Pr(\mathcal{E}) = 1$ only if $\Pr(\mathcal{C}) = 1$.

¹Even though the result there was stated assuming a more restrictive random model, the proof itself relies only on the independence property of the model.




Fig. 2. The ergodicity and consensus coincide *a.s.*, and all three events assume 0-1 law for an i.i.d model.

We next show that the infinite flow event is also a trivial event.

Lemma 4: For an independent random model, the infinite flow event \mathcal{F} occurs with either probability zero or one.

Proof: For a nontrivial set S of $[m]$, the sequence $\{W_S(k)\}$ of undirected flows across the cut (S, \bar{S}) (see Eq. (2)) is a sequence of independent (finitely valued) random variables. The event $\{\sum_{k=0}^{\infty} W_S(k) = \infty\}$ is a tale event and, by Kolmogorov's 0-1 law ([12], page 61), it has either probability 0 or 1. Since there are finitely many nontrivial sets $S \subset [m]$, the event $\mathcal{F} = \bigcap_{S \subset [m]} \{\sum_{k=0}^{\infty} W_S(k) = \infty\}$ also has either probability 0 or 1. ■

While both events \mathcal{E} and \mathcal{F} are trivial for an independent model, the situation is not the same for the consensus event \mathcal{C} . In particular, by Example 3 where $p \in (0, 1)$, we see that the consensus event need not assume 0-1 law since it can occur with a probability $p \in (0, 1)$.

However, the situation is very different for i.i.d. models. In particular, in this case the consensus event is also a trivial event, as seen in the following lemma.

Lemma 5: For an i.i.d. random model, the consensus event \mathcal{C} occurs with either probability zero or one.

Proof: The result follows from the fact that \mathcal{E} is a trivial event and Theorem 2, which states that $\mathcal{E} = \mathcal{C}$ almost surely for i.i.d. models. ■

Figure 2 illustrates the 0-1 laws of \mathcal{E} , \mathcal{C} and \mathcal{F} for an i.i.d. model. Our next example demonstrates that the inclusion relation in Figure 2 can be strict.

Example 4: Consider the independent identical random model where each $W(k)$ is equally likely to be any of the $m \times m$ permutation matrices. Then, in view of the uniform distribution, we have $E[W(k)] = \frac{1}{m} ee^T$ for all k . Hence, by Theorem 3, it follows that the infinite flow event \mathcal{F} is happening almost surely. But, since the chain $\{W(k)\}$ is a sequence of permutation matrices, the consensus event \mathcal{C} never happens. □

C. Random Model and Its Expected Model

Here, we investigate the properties of an independent random model and its corresponding expected model. We establish two results in forthcoming Theorems 3 and 4 that later on play an important role in the establishment the Infinite Flow Theorem in Section V. The first result shows the equivalence of the infinite flow property for a random chain $\{W(k)\}$ and the expected chain $\{E[W(k)]\}$. The result is stated in the following theorem.

Theorem 3: Let $\{W(k)\}$ be an independent random model. Then, the model has infinite flow property if and only if the expected model has infinite flow property, i.e., $\sum_{k=0}^{\infty} E[W_S(k)] = \infty$ for every nontrivial $S \subset [m]$.

Proof: Let $S \subset [m]$ be nontrivial. Since the model is independent, the random variables $W_S(k)$ are independent. By the definition of $W_S(k)$ and the stochasticity of $W(k)$, we have $0 \leq W_S(k) \leq \sum_{i=1}^m \sum_{j=1}^m W_{ij}(k) = m$ for all $k \geq 0$. Thus, by monotone convergence theorem ([12], page 225), the infinite flow property of the model implies the infinite flow property of the expected model. On the other hand, since the model is independent and $0 \leq W_S(k) \leq m$, by Kolmogorov's three-series theorem ([12], page 64) it follows that if $\sum_{k=0}^{\infty} E[W_S(k)] = \infty$, then $\Pr(\sum_{k=0}^{\infty} W_S(k) = \infty) > 0$. Since $\sum_{k=0}^{\infty} W_S(k)$ is a trivial event, we have $\Pr(\sum_{k=0}^{\infty} W_S(k) = \infty) = 1$. This holds for any $S \subset [m]$, implying that the model has infinite flow property. ■

There is no analog of Theorem 3 for the ergodicity or consensus event, unless additional assumptions are imposed. However, a weaker result is available as given in the following lemma.

Lemma 6: Let $\{W(k)\}$ be an independent model and assume that the model admits consensus (is ergodic). Then, the expected model $\{E[W(k)]\}$ reaches a consensus (is ergodic).

The proof² of Lemma 6 can be found in [33], [13].

The following theorem states another important result for later use. In particular, as a consequence of Lemma 6, Theorem 1 and Theorem 3, we establish an equivalent deterministic characterization of the ergodicity event for a class of independent random models.

Theorem 4: Let $\{W(k)\}$ be an independent random model such that $\mathcal{E} = \mathcal{F}$ almost surely. Then, the model is ergodic if and only if the expected model is ergodic.

²Assuming more restrictive assumptions on the model, the result of the lemma was stated in [33] (Theorem 3, (a) \Rightarrow (b)) and [13] (Remark 3.3). However, the proofs in these references rely only on the independence of the model.

Proof: If the ergodicity event is almost sure, then by Lemma 6, the expected model is ergodic. For the converse statement, assume that the expected chain $\{E[W(k)]\}$ is ergodic. Then, by Theorem 1 the chain $\{E[W(k)]\}$ has infinite flow. Therefore, by Theorem 3 the infinite flow event \mathcal{F} is almost sure, and since $\mathcal{E} = \mathcal{F}$ a.s., the ergodicity event is almost sure. ■

IV. MODEL WITH FEEDBACK PROPERTY AND STEADY STATE IN EXPECTATION

In this section, we discuss two properties of a random model that are important in the development of our main results in Section V. In particular, we introduce and study a model with feedback properties and a model with a common steady state in expectation. Stronger forms of these properties have always been used when establishing consensus both for deterministic and random models. Here, we provide some new fundamental insights into these properties.

A. Feedback Properties

We define several types of feedback property. Recall that W^i denotes the i th column vector of a matrix W .

Definition 2: A random model $\{W(k)\}$ has *strong feedback property* if there exists $\gamma > 0$ such that

$$W_{ii}(k) \geq \gamma \quad \text{almost surely for all } k \geq 0 \text{ and all } i \in [m].$$

The model has *feedback property* if there exists $\gamma > 0$ such that

$$E[W_{ii}(k)W_{ij}(k)] \geq \gamma E[W_{ij}(k)] \quad \text{for all } k \geq 0, \text{ and all } i, j \in [m] \text{ with } i \neq j.$$

The model has *weak feedback property* if there exists $\gamma > 0$ such that

$$E[(W^i(k))^T W^j(k)] \geq \gamma (E[W_{ij}(k)] + E[W_{ji}(k)]) \quad \text{for all } k \geq 0, \text{ and all } i, j \in [m] \text{ with } i \neq j.$$

The scalar γ is referred to as a *feedback constant*.

While the difference between feedback and strong feedback property is apparent, the difference between weak feedback and feedback property may not be so obvious. The following example illustrates the difference between these concepts.

Example 5: Consider a static deterministic model $\{A(k)\}$ given by:

$$A(k) = A = \begin{bmatrix} 0 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & 0 & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & 0 \end{bmatrix} \quad \text{for } k \geq 0.$$

Since $A_{ii}A_{ij} = 0$ and $A_{ij} \neq 0$ for all $i \neq j$, the model does not have feedback property. At the same time, since $A_{ij} + A_{ji} = 1$ for $i \neq j$, it follows $(A^i)^T A^j = \frac{1}{4} = \frac{1}{4}(A_{ij} + A_{ji})$. Thus, $\{A(k)\}$ has weak feedback property with $\gamma = \frac{1}{4}$. \square

It can be seen that strong feedback property implies feedback property, which in turn implies weak feedback property. The deterministic consensus and averaging models in [36], [38], [28], [24] deal with the matrices that have non-zero diagonal entries and have non-zero entries uniformly bounded. This requirement is more restrictive than the strong feedback property.

We next establish a relation showing that feedback property of model $\{W(k)\}$ implies strong feedback property for the expected model $\{E[W(k)]\}$.

Lemma 7: Let a random model $\{W(k)\}$ have feedback property with constant γ . Then, the expected model $\{E[W(k)]\}$ has strong feedback property with $\frac{\gamma}{m}$.

Proof: Let the model have feedback property with constant $\gamma > 0$. Then, by the definition of the feedback property, for any time $k \geq 0$ and any $i, j \in [m]$ with $i \neq j$, we have $E[W_{ii}(k)W_{ij}(k)] \geq \gamma E[W_{ij}(k)]$. Since $W_{ij}(k) \leq 1$, it follows that

$$E[W_{ii}(k)] \geq E[W_{ii}(k)W_{ij}(k)] \geq \gamma E[W_{ij}(k)] \quad \text{for all } i, j \in [m], i \neq j.$$

The matrices $W(k)$ are stochastic, so that we have $\sum_{j=1}^m E[W_{ij}(k)] = 1$. Hence, for every $k \geq 0$ and $i \in [m]$, there exists an index j^* (the dependence on k and i is suppressed) such that $E[W_{ij^*}(k)] \geq \frac{1}{m}$. If $j^* = i$, then we are done; otherwise we have $E[W_{ii}(k)] \geq \gamma E[W_{ij^*}(k)] \geq \frac{\gamma}{m}$ for all $i \in [m]$. Hence, the expected chain has the strong feedback property with constant $\frac{\gamma}{m}$. \blacksquare

We now focus on an independent model. We have the following result.

Lemma 8: Consider an independent model $\{W(k)\}$. Suppose that the model is such that there is an $\eta > 0$ with the following property: for all $k \geq 0$ and $i, j \in [m]$ with $i \neq j$,

$$E[W_{ij}(k)] > 0 \implies E[W_{ii}(k)W_{ij}(k)] \geq \eta,$$

or the following property: for all $k \geq 0$ and $i, j \in [m]$ with $i \neq j$,

$$E[W_{ij}(k)] > 0 \implies E[(W^i(k))^T W^j(k)] \geq \eta.$$

Then, respectively, the model has feedback property with constant η or weak feedback property with constant $\eta/2$.

Proof: We prove only the case of feedback property, since the other case uses the same line of argument. If $E[W_{ij}(k)] = 0$ for some k and i, j with $i \neq j$, then the relation $E[W_{ii}(k)W_{ij}(k)] \geq$

$\eta \mathbb{E}[W_{ij}(k)]$ is satisfied trivially (with both sides equal to zero). If $\mathbb{E}[W_{ij}(k)] > 0$, then by the assumption of the lemma, we have $\mathbb{E}[W_{ij}(k)W_{ii}(k)] \geq \eta$. Furthermore, since $1 \geq W_{ij}(k)$ for all i, j and k , it follows that $\mathbb{E}[W_{ij}(k)W_{ii}(k)] \geq \eta \geq \eta \mathbb{E}[W_{ij}(k)]$, thus showing that the model has feedback property with constant η . ■

The i.i.d. models $\{W(k)\}$ with almost surely positive diagonal entries $W_{ii}(k)$ have been studied in [13], [33], [34]. These models have feedback property, as seen in the following corollary.

Corollary 1: Let $\{W(k)\}$ be an i.i.d. model with almost surely positive diagonal entries. Then, the model has feedback property with constant $\gamma = \min_{\{i \neq j | \mathbb{E}[W_{ij}(k)] > 0\}} \mathbb{E}[W_{ii}(k)W_{ij}(k)]$.

Proof: Since $W_{ii}(k) > 0$ a.s., it follows $W_{ii}(k)W_{ij}(k)1_{\{W_{ij}(k) > 0\}} > 0$ a.s., implying that $\mathbb{E}[W_{ii}(k)W_{ij}(k)] > 0$ whenever $\mathbb{E}[W_{ij}(k)] > 0$. Define $\eta = \min_{\{i \neq j | \mathbb{E}[W_{ij}(k)] > 0\}} \mathbb{E}[W_{ii}(k)W_{ij}(k)]$. Since the model is i.i.d., the constant η is independent of time. Hence, by Lemma 8 it follows that the model has feedback property with constant η . ■

B. Steady State in Expectation

Here, we consider a model with another special property. Specifically, we discuss a random model $\{W(k)\}$ such that its expected chain $\{\mathbb{E}[W(k)]\}$ has a common steady state.

Definition 3: A random model has a *common steady state in expectation* if there is a stochastic vector $\pi \in \mathbb{R}^m$ such that $\pi^T \mathbb{E}[W(k)] = \pi^T$ for all k .

For example, the $m \times m$ matrices that are doubly stochastic in expectation satisfy the preceding definition with $\pi = \frac{1}{m} \mathbf{e}$, such as the matrices arising in a randomized broadcast or gossip over a connected (static) network [1], [6].

Consider the function given by

$$V(x) = \sum_{i=1}^m \pi_i (x_i - \pi^T x)^2 \quad \text{for } x \in \mathbb{R}^m. \quad (6)$$

The function $V(x)$ measures the weighted spread of the vector x entries with respect to the weighted average value $\pi^T x$. It can also be viewed as a weighted distance between x and $\pi^T x \mathbf{e}$.

We at first study the behavior of the weighted averages $\pi^T x(k)$ along the sequence $\{x(k)\}$ starting with any initial state $x(0) \in \mathbb{R}^m$. The main observation is that the random scalar sequence $\{\pi^T x(k)\}$ is a bounded martingale, which leads us to the following result.

Lemma 9: Let the random model be independent with a common steady state π in expectation. Then, the sequence $\{\pi^T x(k)\}$ converges almost surely for any $x(0) \in \mathbb{R}^m$.

Proof: By the independence of the model and the property $\pi^T \mathbb{E}[W(k)] = \pi^T$, it follows that, for any initial $x(0) \in \mathbb{R}^m$, the process $\{\pi^T x(k)\}$ is a martingale with respect to the natural filtration of the process. Since the matrices $W(k)$ are stochastic, the sequence $\{x(k)\}$ is bounded, thus implying that $\{\pi^T x(k)\}$ is a bounded martingale. By the martingale convergence theorem (see [3], Theorem 35.5), the sequence $\{\pi^T x(k)\}$ converges almost surely. ■

We next characterize the limiting value of the martingale $\{\pi^T x(k)\}$. Let $\lambda_{i,\pi} \in \mathbb{R}$ be the limit of the martingale $\{\pi^T x(k)\}$ for the initial state $x(0) = e_i$, and let λ_π be the vector defined by

$$\lambda_\pi = (\lambda_{1,\pi}, \dots, \lambda_{m,\pi}) \quad \text{with} \quad \lambda_{i,\pi} = \lim_{k \rightarrow \infty} \pi^T W(k) \cdots W(0) e_i \quad \text{for } i \in [m]. \quad (7)$$

In the following theorem, we provide some properties of the random vector λ_π .

Lemma 10: Let the random model be independent with a common steady state π in expectation. Then, the random vector λ_π has the following properties:

- (a) $\lim_{k \rightarrow \infty} \pi^T x(k) = \lambda_\pi^T x(0)$ almost surely for any $x(0) \in \mathbb{R}^m$.
- (b) λ_π is a stochastic vector, i.e., $\lambda_\pi \geq 0$ and $\lambda_\pi^T e = 1$.
- (c) $\mathbb{E}[\lambda_\pi] = \pi$.
- (d) For a given $x(0) \in \mathbb{R}^m$, the limit points of the sequence $\{x(k)\}$ lie in the random hyperplane $\mathcal{H}_{\pi, \lambda_\pi^T x(0)} = \{x \in \mathbb{R}^m \mid \pi^T x = \lambda_\pi^T x(0)\}$ almost surely.

Proof: By Lemma 9 and the definition of $\lambda_{i,\pi}$, we have $\lim_{k \rightarrow \infty} \pi^T x(k) = \lambda_{i,\pi}$ almost surely for initial state e_i . Using the linearity of the system in (1), we obtain for any $x(0) \in \mathbb{R}^m$,

$$\lim_{k \rightarrow \infty} \pi^T x(k) = \sum_{i=1}^m x_i(0) \left(\lim_{k \rightarrow \infty} \pi^T W(k-1) \cdots W(0) e_i \right) = \sum_{i=1}^m x_i(0) \lambda_{i,\pi} = \lambda_\pi^T x(0),$$

thus showing part (a).

Since $x(k) = W(k-1) \cdots W(0)x(0)$, by part (a) it follows $\lim_{k \rightarrow \infty} \pi^T W(k-1) \cdots W(0) = \lambda_\pi^T$. The matrices $W(k)$ and the vector π have nonnegative entries implying that the vector λ_π also has nonnegative entries. By letting $x(0) = e$ and using the stochasticity of $W(k)$, we have $x(k) = e$ for all k , implying $1 = \pi^T x(k)$ for all k . Thus, we have $1 = \lim_{k \rightarrow \infty} \pi^T x(k) = \lambda_\pi^T x(0) = \lambda_\pi^T e$, where the second equality holds by part (a). Hence, λ_π is a stochastic vector.

To show part (c), we note that by the martingale property of the process $\{\pi^T x(k)\}$, we have

$$\mathbb{E}[\pi^T x(k)] = \pi^T x(0) \quad \text{for all } k \geq 0 \text{ and any } x(0) \in \mathbb{R}^m.$$

By the boundedness of the martingale for any $x(0) \in \mathbb{R}^m$, we have

$$\lim_{k \rightarrow \infty} \mathbb{E}[\pi^T x(k)] = \mathbb{E}[\lambda_\pi^T] x(0) \quad \text{for any } x(0) \in \mathbb{R}^m.$$

The preceding two relations imply $E[\lambda_\pi] = \pi$.

For the part (d), we note that by the stochasticity of $W(k)$, the sequence $\{x(k)\}$ is bounded for every initial state $x(0)$. Thus, $\{x(k)\}$ has some limit points. By part (a), each limit point x^* of the sequence satisfies $\pi^T x^* = \lambda_\pi^T x(0)$ almost surely. ■

We now focus on the sequence $\{V(x(k))\}$ and we show that it is a convergent supermartingale. In fact, this indicates that $V(x)$ is a stochastic Lyapunov function for the random system in (1).

Theorem 5: Let the random model be independent with a common steady state π in expectation. Then, we almost surely have for all $k \geq 0$,

$$E[V(x(k+1)) \mid x(k)] \leq V(x(k)) - \sum_{i < j} H_{ij}(k) (x_i(k) - x_j(k))^2,$$

where $H(k) = E[W^T(k)DW(k)]$. Furthermore, $\{V(x(k))\}$ converges almost surely.

Proof: By using $D = \text{diag}(\pi)$, from the definition of the function $V(x)$ in (6) we have

$$V(x) = x^T(I - \pi e^T)D(I - e\pi^T)x = x^T(D - \pi\pi^T)x,$$

where the second equality is obtained by using $e^T D = \pi^T$, $De = \pi$, and $\pi^T e = 1$. In view of $x(k+1) = W(k)x(k)$, it follows that for all $k \geq 0$,

$$V(x(k+1)) = x(k+1)^T(D - \pi\pi^T)x(k+1) = x(k)W(k)^T(D - \pi\pi^T)W(k)x(k). \quad (8)$$

Since the model is independent, by taking the expectation conditioned on $x(k)$, we obtain almost surely for all $k \geq 0$,

$$E[V(x(k+1)) \mid x(k)] = x(k)^T E[W(k)^T(D - \pi\pi^T)W(k)] x(k).$$

Since $H(k) = E[W^T(k)DW(k)]$, we further have

$$\begin{aligned} E[V(x(k+1)) \mid x(k)] &= x(k)^T H(k)x(k) - E\left[\left(\pi^T W(k)x(k)\right)^2 \mid x(k)\right] \\ &\leq x(k)^T H(k)x(k) - \left(E\left[\pi^T W(k)x(k) \mid x(k)\right]\right)^2, \end{aligned}$$

where the inequality follows by Jensen's inequality (see [12], page 225) and the convexity of the function $s \mapsto s^2$. The expected matrices $E[W(k)]$ have the same steady state π , implying that almost surely

$$E\left[\pi^T W(k)x(k) \mid x(k)\right] = \pi^T E[W(k)]x(k) = \pi^T x(k) \quad \text{for all } k \geq 0.$$

By combining the preceding two relations, we see that almost surely for all $k \geq 0$,

$$\mathbb{E}[V(x(k+1)) \mid x(k)] \leq x(k)^T (H(k) - \pi\pi^T) x(k).$$

By adding and subtracting $x(k)^T D x(k)$ to the right hand side of the preceding relation and using (8), we obtain almost surely

$$\mathbb{E}[V(x(k+1)) \mid x(k)] \leq x(k)^T (H(k) - D) x(k) + V(x(k)) \quad \text{for all } k \geq 0. \quad (9)$$

We now show that $x(k)^T (H(k) - D) x(k) = \sum_{i < j} H_{ij}(k)(x_i(k) - x_j(k))^2$. By the definition of $H(k)$, we have $H_{ij}(k) = \sum_{\ell=1}^m \mathbb{E}[\pi_\ell W_{\ell i}(k) W_{\ell j}(k)]$, so that for any $i \in [m]$,

$$\sum_{\substack{j=1 \\ j \neq i}}^m H_{ij}(k) = \mathbb{E} \left[\sum_{\substack{j=1 \\ j \neq i}}^m \sum_{\ell=1}^m \pi_\ell W_{\ell i}(k) W_{\ell j}(k) \right] = \mathbb{E} \left[\sum_{\ell=1}^m \pi_\ell W_{\ell i}(k) \sum_{\substack{j=1 \\ j \neq i}}^m W_{\ell j}(k) \right].$$

Since $W(k)$ is stochastic, we have $\sum_{j=1, j \neq i}^m W_{\ell j}(k) = 1 - W_{\ell i}(k)$, implying that

$$\sum_{\substack{j=1 \\ j \neq i}}^m H_{ij}(k) = \mathbb{E} \left[\sum_{\ell=1}^m \pi_\ell W_{\ell i}(k) \right] - \mathbb{E} \left[\sum_{\ell=1}^m \pi_\ell W_{\ell i}^2(k) \right] = \pi_i - H_{ii}(k),$$

where the last equality follows from $\mathbb{E}[\pi^T W(k)] = \pi^T$. Since $D = \text{diag}(\pi)$, the preceding relation yields

$$H_{ii}(k) = D_{ii} - \sum_{\substack{j=1 \\ j \neq i}}^m H_{ij}(k). \quad (10)$$

Therefore, for any $x \in \mathbb{R}^m$, we have $x^T H(k) x = \sum_{i=1}^m x_i \sum_{j=1}^m H_{ij}(k) x_j$, which can be further written as

$$x^T H(k) x = \sum_{i=1}^m x_i \sum_{\substack{j=1 \\ j \neq i}}^m H_{ij}(k) x_j + \sum_{i=1}^m x_i H_{ii}(k) x_i = \sum_{i=1}^m x_i \sum_{\substack{j=1 \\ j \neq i}}^m H_{ij}(k) (x_j - x_i) + \sum_{i=1}^m x_i D_{ii} x_i,$$

where the last equality follows from relation (10). The matrix $H(k) = \mathbb{E}[W(k)^T D W(k)]$ is symmetric, so that

$$\sum_{i=1}^m x_i \sum_{\substack{j=1 \\ j \neq i}}^m H_{ij}(k) (x_j - x_i) = - \sum_{i < j} H_{ij}(k) (x_i - x_j)^2,$$

implying that $x^T H(k) x = - \sum_{i < j} H_{ij}(k) (x_i - x_j)^2 + x^T D x$. Therefore, we have

$$x(k)^T (H(k) - D) x(k) = - \sum_{i < j} H_{ij}(k) (x_i(k) - x_j(k))^2.$$

By combining the preceding relation with (9), we obtain almost surely

$$\mathbb{E}[V(x(k+1)) \mid x(k)] \leq V(x(k)) - \sum_{i < j} H_{ij}(k) (x_i(k) - x_j(k))^2 \quad \text{for all } k \geq 0.$$

Since $W(k)$ are stochastic matrices and π is stochastic vector, the matrix $H(k)$ has non-negative entries for all k . Hence, from the preceding relation it follows that $\{V(x(k))\}$ is a supermartingale. The convergence of $\{V(x(k))\}$ follows straightforwardly from the nonnegative supermartingale convergence (see [12], (2.11) Corollary, page 236.) ■

We conclude this section with another result for the weighted distance function $V(x)$. This result plays crucial role in establishing our result in Section V-B.

Lemma 11: Let $\pi \in \mathbb{R}^m$ be a stochastic vector, and let $x \in \mathbb{R}^m$ be such that $x_1 \leq \dots \leq x_m$. Then, for $V(x)$ we have

$$\frac{1}{(m-1)^2} V(x) \leq \frac{1}{x_m - x_1} \sum_{i=1}^{m-1} (x_{i+1} - x_i)^3.$$

Proof: We establish two relations

$$\frac{1}{m-1} V(x) \leq \sum_{i=1}^{m-1} (x_{i+1} - x_i)^2, \quad (11)$$

$$\frac{1}{m-1} \sum_{i=1}^{m-1} (x_{i+1} - x_i)^2 \leq \frac{1}{x_m - x_1} \sum_{i=1}^{m-1} (x_{i+1} - x_i)^3. \quad (12)$$

Observe that the desired inequality follows from relations (11)–(12).

We now show relation (11). We have $x_i \leq x_m$ for all i . Since π is stochastic, we also have $\pi^T x \geq x_1$. Thus,

$$V(x) = \sum_{i=1}^m \pi_i (x_i - \pi^T x)^2 \leq \sum_{i=1}^m \pi_i (x_m - x_1)^2 = (x_m - x_1)^2.$$

By writing $x_m - x_1 = \sum_{i=1}^{m-1} (x_{i+1} - x_i) = (m-1) \left(\frac{1}{m-1} \sum_{i=1}^{m-1} (x_{i+1} - x_i) \right)$, we obtain

$$(x_m - x_1)^2 = (m-1)^2 \left(\frac{1}{m-1} \sum_{i=1}^{m-1} (x_{i+1} - x_i) \right)^2 \leq (m-1) \sum_{i=1}^{m-1} (x_{i+1} - x_i)^2,$$

where the last inequality holds by the convexity of the function $s \mapsto s^2$. The preceding two relations yield $V(x) \leq (m-1) \sum_{i=1}^{m-1} (x_{i+1} - x_i)^2$, thus implying relation (11).

To prove relation (12), we write

$$\begin{aligned} (x_m - x_1) \sum_{i=1}^{m-1} (x_{i+1} - x_i)^2 &= \sum_{j=1}^{m-1} (x_{j+1} - x_j) \sum_{i=1}^{m-1} (x_{i+1} - x_i)^2 \\ &= \sum_{j=1}^{m-1} (x_{j+1} - x_j)^3 + \sum_{j < i} \left((x_{j+1} - x_j)(x_{i+1} - x_i)^2 + (x_{i+1} - x_i)(x_{j+1} - x_j)^2 \right). \end{aligned}$$

To estimate the last term in the preceding relation, we consider scalars $\alpha \geq 0$ and $\beta \geq 0$, and let $u = (\alpha, \beta)$ and $v = (\beta^2, \alpha^2)$. Then, by Hölder's inequality with $p = 3$, $q = \frac{3}{2}$, we have $u^T v \leq \|u\|_p \|v\|_q$, where $\|\cdot\|_p$ is the p -norm. Hence,

$$\alpha\beta^2 + \beta\alpha^2 \leq (\alpha^3 + \beta^3)^{\frac{1}{3}} (\beta^3 + \alpha^3)^{\frac{2}{3}} = \alpha^3 + \beta^3. \quad (13)$$

By using inequality (13) with $\alpha_j = (x_{j+1} - x_j)$ and $\beta_i = (x_{i+1} - x_i)$ for different indices j and i , $1 \leq j < i \leq m-1$, we obtain

$$\begin{aligned} (x_m - x_1) \sum_{i=1}^{m-1} (x_{i+1} - x_i)^2 &\leq \sum_{j=1}^{m-1} (x_{j+1} - x_j)^3 + \sum_{j < i} \left((x_{j+1} - x_j)^3 + (x_{i+1} - x_i)^3 \right) \\ &= (m-1) \sum_{i=1}^{m-1} (x_{i+1} - x_i)^3. \end{aligned}$$

Relation (12) now follows by dividing with $m-1$. ■

V. MODEL WITH INFINITE FLOW PROPERTY

We consider an independent random model with infinite flow. We show that this property, together with weak feedback and a common steady state in expectation, is necessary and sufficient for almost sure ergodicity. Moreover, we establish that the ergodicity of the model is equivalent to the ergodicity of the expected model.

A. Preliminary Result

We now provide an important relation that we use later on in Section V-B.

Lemma 12: Let $\{A(k)\} \subset \mathbb{S}^m$ and $z(k+1) = A(k)z(k)$ for all $k \geq 0$, starting with some $z(0) \in \mathbb{R}^m$. Let σ be a permutation of the index set $[m]$ corresponding to the nondecreasing ordering of the entries $z_\ell(0)$, i.e., σ is a permutation on $[m]$ such that $z_{\sigma_1}(0) \leq \dots \leq z_{\sigma_m}(0)$. Also, let $T \geq 1$ be such that

$$\sum_{k=0}^{T-1} A_S(k) \geq \delta \quad \text{for every nontrivial set } S \subset [m], \quad (14)$$

where $\delta \in (0, 1)$ is arbitrary. Then, we have

$$\sum_{k=0}^{T-1} \sum_{i < j} (A_{ij}(k) + A_{ji}(k)) (z_j(k) - z_i(k))^2 \geq \frac{\delta(1-\delta)^2}{z_{\sigma_m}(0) - z_{\sigma_1}(0)} \sum_{i=1}^{m-1} (z_{\sigma_{i+1}}(0) - z_{\sigma_i}(0))^3.$$

Proof: Since relation (14) holds for any nontrivial set $S \subset [m]$, without loss of generality we may assume that the permutation σ is identity (otherwise we will relabel the indices of the entries in $z(0)$ and update the matrices accordingly). Hence, we have $z_1(0) \leq \dots \leq z_m(0)$. For each $\ell = 1, \dots, m-1$, let $S_\ell = \{1, \dots, \ell\}$ and define time $t_\ell \geq 1$, as follows:

$$t_\ell = \operatorname{argmin}_{t \geq 1} \left\{ \sum_{k=0}^{t-1} A_{S_\ell}(k) \geq \delta \frac{z_{\ell+1}(0) - z_\ell(0)}{z_m(0) - z_1(0)} \right\}.$$

Since the entries of $z(0)$ are nondecreasing, we have $\delta \frac{z_{\ell+1}(0) - z_\ell(0)}{(z_m(0) - z_1(0))} \leq \delta$ for all $\ell = 1, \dots, m-1$. Thus, by relation (14), the time $t_\ell \geq 1$ exists and $t_\ell \leq T$ for each ℓ .

We next estimate $z_j(k) - z_i(k)$ for all $i < j$ and any time $k = 0, \dots, T-1$. For this, we introduce the following index sets $a_{ij}(k) \subset [m]$,

$$a_{ij}(k) = \{\ell \in [m] \mid k \leq t_\ell - 1, \ell \geq i, \ell + 1 \leq j\} \quad \text{for } 0 \leq k \leq T-1, \text{ and } i < j.$$

Let $k \leq t_\ell - 1$ for some ℓ . Since $S_\ell = \{1, \dots, \ell\}$, we have $i \in S_\ell$ and $j \in \bar{S}_\ell$. Thus, by Lemma 1 we have for any $k \geq 1$,

$$\begin{aligned} z_i(k) &\leq \max_{s \in S_\ell} z_s(0) + (z_m(0) - z_1(0)) \sum_{\tau=0}^{k-1} A_{S_\ell}(\tau), \\ z_j(k) &\geq \min_{r \in \bar{S}_\ell} z_r(0) - (z_m(0) - z_1(0)) \sum_{\tau=0}^{k-1} A_{S_\ell}(\tau). \end{aligned}$$

Furthermore, we have $\max_{s \in S_\ell} z_s(0) = z_\ell(0)$ and $\min_{r \in \bar{S}_\ell} z_r(0) = z_{\ell+1}(0)$ since $S_\ell = \{1, \dots, \ell\}$ and $z_1(0) \leq \dots \leq z_m(0)$. Thus, it follows

$$\begin{aligned} z_i(k) - z_\ell(0) &\leq (z_m(0) - z_1(0)) \sum_{\tau=0}^{k-1} A_{S_\ell}(\tau), \\ z_{\ell+1}(0) - z_j(k) &\leq (z_m(0) - z_1(0)) \sum_{\tau=0}^{k-1} A_{S_\ell}(\tau). \end{aligned}$$

By the definition of time t_ℓ , we have $(z_m(0) - z_1(0)) \sum_{\tau=0}^{t_\ell-1} A_{S_\ell}(\tau) < \delta (z_{\ell+1}(0) - z_\ell(0))$ for $k \leq t_\ell - 1$. Hence, by using this and the definition of $a_{ij}(k)$, we have

$$z_i(k) - z_\ell(0) \leq \delta (z_{\ell+1}(0) - z_\ell(0)) \quad \text{for any } \ell \in a_{ij}(k), \quad (15)$$

$$z_{\ell+1}(0) - z_j(k) \leq \delta(z_{\ell+1}(0) - z_\ell(0)) \quad \text{for any } \ell \in a_{ij}(k). \quad (16)$$

Now suppose that $a_{ij}(k) = \{\ell_1, \dots, \ell_r\}$ for some $r \leq m-1$, and assume that $\ell_1 \leq \dots \leq \ell_r$. By choosing $\ell = \ell_1$ in (15) and $\ell = \ell_r$ in (16), and by letting $\alpha_i = z_{i+1}(0) - z_i(0)$, we obtain

$$z_j(k) - z_i(k) \geq z_{\ell_r+1}(0) - z_{\ell_1}(0) - \delta(\alpha_{\ell_r} + \alpha_{\ell_1}).$$

Since $z_i(0) \leq z_{i+1}(0)$ for all $i = 1, \dots, m-1$, we have

$$z_{\ell_1}(0) \leq z_{\ell_1+1}(0) \leq z_{\ell_2}(0) \leq z_{\ell_2+1}(0) \leq \dots \leq z_{\ell_r}(0) \leq z_{\ell_r+1}(0),$$

which combined with the preceding relation yields

$$z_j(k) - z_i(k) \geq \sum_{\xi=1}^{r-1} (z_{\ell_\xi+1}(0) - z_{\ell_\xi}(0)) - \delta(\alpha_{\ell_r} + \alpha_{\ell_1}).$$

Using $\alpha_i = z_{i+1}(0) - z_i(0)$ and $a_{ij}(k) = \{\ell_1, \dots, \ell_r\}$, we further have

$$z_j(k) - z_i(k) \geq \sum_{\xi=1}^{r-1} \alpha_{\ell_\xi} - \delta(\alpha_{\ell_r} + \alpha_{\ell_1}) \geq (1-\delta) \sum_{\xi=1}^{r-1} \alpha_{\ell_\xi} = (1-\delta) \sum_{\ell \in a_{ij}(k)} \alpha_\ell.$$

From the preceding relation it follows that

$$\begin{aligned} \sum_{i < j} (A_{ij}(k) + A_{ji}(k)) (z_j(k) - z_i(k))^2 &\geq (1-\delta)^2 \sum_{i < j} (A_{ij}(k) + A_{ji}(k)) \left(\sum_{\ell \in a_{ij}(k)} \alpha_\ell \right)^2 \\ &\geq (1-\delta)^2 \sum_{i < j} (A_{ij}(k) + A_{ji}(k)) \left(\sum_{\ell \in a_{ij}(k)} \alpha_\ell^2 \right), \end{aligned}$$

where the last inequality holds by $\alpha_\ell \geq 0$. In the last term in the preceding relation, the coefficient of α_ℓ^2 is equal to $(1-\delta)^2 A_{S_\ell}(k)$. Furthermore, by the definition of $a_{ij}(k)$, we have $\ell \in a_{ij}(k)$ only when $k \leq t_\ell - 1$. Therefore,

$$\begin{aligned} \sum_{i < j} (A_{ij}(k) + A_{ji}(k)) (z_j(k) - z_i(k))^2 &\geq (1-\delta)^2 \sum_{i < j} (A_{ij}(k) + A_{ji}(k)) \left(\sum_{\ell \in a_{ij}(k)} \alpha_\ell^2 \right) \\ &= (1-\delta)^2 \sum_{\{\ell | k \leq t_\ell - 1\}} A_{S_\ell}(k) \alpha_\ell^2. \end{aligned}$$

By summing the preceding relations over $k = 0, \dots, T-1$, we obtain

$$\begin{aligned} \sum_{k=0}^{T-1} \sum_{i < j} (A_{ij}(k) + A_{ji}(k)) (z_j(k) - z_i(k))^2 &\geq (1-\delta)^2 \sum_{k=0}^{T-1} \sum_{\{\ell | k \leq t_\ell - 1\}} A_{S_\ell}(k) \alpha_\ell^2 \\ &\geq (1-\delta)^2 \sum_{\ell=1}^{m-1} \left(\sum_{k=0}^{t_\ell-1} A_{S_\ell}(k) \right) \alpha_\ell^2, \end{aligned}$$

where the last inequality follows by exchanging the order of summation. By the definition of t_ℓ and using $\alpha_\ell = z_{\ell+1}(0) - z_\ell(0)$, we have $\sum_{k=0}^{t_\ell-1} A_{S_\ell}(k) \geq \delta \alpha_\ell / (z_m(0) - z_1(0))$, implying

$$\begin{aligned} \sum_{k=0}^{T-1} \sum_{i < j} (A_{ij}(k) + A_{ji}(k)) (z_j(k) - z_i(k))^2 &\geq (1-\delta)^2 \sum_{\ell=1}^{m-1} \delta \frac{\alpha_\ell}{z_m(0) - z_1(0)} \alpha_\ell^2 \\ &= \delta(1-\delta)^2 \sum_{\ell=1}^{m-1} \frac{\alpha_\ell^3}{z_m(0) - z_1(0)}. \end{aligned}$$

■

B. Sufficient Conditions for Ergodicity

We now establish one of our main results pertinent to independent random models with infinite flow property and having a common vector in expectation and weak feedback property.

Let $t_0 = 0$ and for any $q \geq 1$, let

$$t_q = \operatorname{argmin}_{t \geq t_{q-1}+1} \Pr \left(\min_{S \subset [m]} \sum_{k=t_{q-1}}^{t-1} W_S(k) \geq \delta \right) \geq \epsilon, \quad (17)$$

where $\epsilon, \delta \in (0, 1)$ are arbitrary. For each $q \geq 0$, define

$$\mathcal{A}_q = \left\{ \omega \mid \min_{S \subset [m]} \sum_{k=t_q}^{t_{q+1}-1} W_S(k)(\omega) \geq \delta \right\}. \quad (18)$$

Since the model has infinite flow property, the infinite flow event \mathcal{F} occurs almost surely. Therefore, the time t_q is finite for all q , i.e. $t_q < \infty$.

The following result shows that either the infinite flow or expected infinite flow property is sufficient for the ergodicity of the model.

Theorem 6: (Sufficient Ergodicity Condition) Let $\{W(k)\}$ be an independent random model with a common steady state $\pi > 0$ in expectation and weak feedback property. Also, let the model have either infinite flow or expected infinite flow property. Then, the model is ergodic. In

particular, $\lim_{k \rightarrow \infty} x_i(k) = \lambda_\pi^T x(0)$ almost surely for all $i \in [m]$, where λ_π is the random vector of Eq. (7).

Proof: Consider the case when the model has infinite flow property. Let $x(0) \in \mathbb{R}^m$ be arbitrary initial state. Let us denote the (random) ordering of the entries of the vector $x(t_q)$ by η^q for all q . Thus, at time t_q , we have $x_{\eta_1^q}(t_q) \leq \dots \leq x_{\eta_m^q}(t_q)$.

Now, let $q \geq 0$ be arbitrary and fixed, and consider the set \mathcal{A}_q in (18). By the definition of \mathcal{A}_q , we have $\sum_{k=t_q}^{t_{q+1}-1} W_S(k)(\omega) \geq \delta$ for any $S \subset [m]$ and $\omega \in \mathcal{A}_q$. Thus, by Lemma 12, we obtain for any $\omega \in \mathcal{A}_q$,

$$\sum_{t=t_q}^{t_{q+1}-1} \sum_{i < j} (W_{ij}(t) + W_{ji}(t)) (x_i(t) - x_j(t))^2(\omega) \geq \frac{\delta(1-\delta)^2}{d(t_q)(\omega)} \sum_{\ell=1}^{m-1} (x_{\eta_{\ell+1}^q}(t_q) - x_{\eta_\ell^q}(t_q))^3(\omega),$$

where $d(t_q) = x_{\eta_m^q}(t_q) - x_{\eta_1^q}(t_q)$. On the other hand, by Lemma 11, we have

$$\frac{1}{x_{\eta_m^q}(t_q)(\omega) - x_{\eta_1^q}(t_q)(\omega)} \sum_{\ell=1}^{m-1} (x_{\eta_{\ell+1}^q}(t_q) - x_{\eta_\ell^q}(t_q))^3(\omega) \geq \frac{1}{(m-1)^2} V(x(t_q))(\omega).$$

Hence, for any $\omega \in \mathcal{A}_q$,

$$\sum_{t=t_q}^{t_{q+1}-1} \sum_{i < j} (W_{ij}(t) + W_{ji}(t)) (x_i(t) - x_j(t))^2(\omega) \geq \frac{\delta(1-\delta)^2}{(m-1)^2} V(x(t_q))(\omega),$$

which we can compactly write as follows

$$\sum_{t=t_q}^{t_{q+1}-1} \sum_{i < j} (W_{ij}(t) + W_{ji}(t)) (x_i(t) - x_j(t))^2 \geq \frac{\delta(1-\delta)^2}{(m-1)^2} V(x(t_q)) 1_{\mathcal{A}_q}, \quad (19)$$

where $1_{\mathcal{A}_q}$ is the indicator function of the event \mathcal{A}_q .

Observe that $x(t)$ and $W(t)$ are independent since the model is independent. Therefore, by Theorem 5, we have

$$\mathbb{E}[V(x(t_{q+1}))] - \mathbb{E}[V(x(t_q))] \leq - \sum_{t=t_q}^{t_{q+1}-1} \sum_{i < j} H_{ij}(t) \mathbb{E}[(x_i(t) - x_j(t))^2],$$

with $H(k) = \mathbb{E}[W^T(k)DW(k)]$ and $D = \text{diag}(\pi)$. Let $\pi_{\min} = \min_{\ell \in [m]} \pi_\ell$ and note that $\pi_{\min} > 0$ since $\pi > 0$. Thus, we have

$$H_{ij}(t) = \mathbb{E}[(W^i(t))^T DW^j(t)] \geq \pi_{\min} \mathbb{E}[(W^i(t))^T W^j(t)] \geq \pi_{\min} \gamma (\mathbb{E}[W_{ij}(k)] + \mathbb{E}[W_{ji}(k)]).$$

Therefore,

$$\mathbb{E}[V(x(t_{q+1}))] - \mathbb{E}[V(x(t_q))] \leq -\pi_{\min} \gamma \mathbb{E} \left[\sum_{t=t_q}^{t_{q+1}-1} \sum_{i < j} (W_{ij}(k) + W_{ji}(k)) (x_i(t) - x_j(t))^2 \right].$$

Further, by using relation (19), we obtain

$$\begin{aligned} \mathbb{E}[V(x(t_{q+1}))] - \mathbb{E}[V(x(t_q))] &\leq -\pi_{\min} \gamma \mathbb{E}\left[\frac{\delta(1-\delta)^2}{(m-1)^2} 1_{A_q} V(x(t_q))\right] \\ &= -\frac{\delta(1-\delta)^2 \gamma \pi_{\min}}{(m-1)^2} \mathbb{E}[1_{A_q} V(x(t_q))] \\ &\leq -\frac{\epsilon \delta(1-\delta)^2 \gamma \pi_{\min}}{(m-1)^2} \mathbb{E}[V(x(t_q))], \end{aligned}$$

where the last inequality follows by $\Pr(\mathcal{A}_q) \geq \epsilon$, and the fact that $1_{\mathcal{A}_q}$ and $V(x(t_q))$ are independent (since $x(t_q)$ depends on information prior to time t_q and the set \mathcal{A}_q relies on information at time t_q and later). Hence, it follows

$$\mathbb{E}[V(x(t_{q+1}))] \leq \left(1 - \frac{\epsilon \delta(1-\delta)^2 \gamma \pi_{\min}}{(m-1)^2}\right) \mathbb{E}[V(x(t_q))].$$

Therefore, since $q \geq 0$ is arbitrary, we see

$$\mathbb{E}[V(x(t_q))] \leq \left(1 - \frac{\epsilon \delta(1-\delta)^2 \gamma \pi_{\min}}{(m-1)^2}\right)^q \mathbb{E}[V(x(0))] \quad \text{for all } q \geq 0,$$

implying that $\sum_{q=0}^{\infty} \mathbb{E}[V(x(t_q))] < \infty$. In view of the nonnegativity of $V(x)$, by the monotone convergence theorem ([3], Theorem 16.6) it follows $\mathbb{E}\left[\sum_{q=0}^{\infty} V(x(t_q))\right] < \infty$, implying $\lim_{q \rightarrow \infty} V(x(t_q)) = 0$ *a.s.* According to Theorem 5, we have that the sequence $\{V(x(k))\}$ is convergent, which together with the preceding relation implies that $\lim_{k \rightarrow \infty} V(x(k)) = 0$ *a.s.*

To show the ergodicity of the model, we note that by the convergence result for the martingale $\{\pi^T x(k)\}$ in Lemma 10(a), we have $\lim_{k \rightarrow \infty} \pi^T x(k) = \lambda_{\pi}^T x(0)$ almost surely, where the random vector λ_{π} is given by Eq. (7). Now, recalling $V(x) = \sum_{i=1}^m \pi_i (x - \pi^T x)^2$, and using $\pi^T x(k) \rightarrow \lambda_{\pi}^T x(0)$ and the fact that all norms in \mathbb{R}^m are equivalent, we obtain $\lim_{k \rightarrow \infty} |x_i(k) - \lambda_{\pi}^T x(0)| = 0$ *a.s.* for all $i \in [m]$, thus showing the ergodicity of the model.

Consider now the case when the model has expected infinite flow, i.e., $\sum_{k=0}^{\infty} \mathbb{E}[W_S(k)] = \infty$ for every nontrivial $S \subset [m]$. By Theorem 3, the model $\{W(k)\}$ has infinite flow property if and only if it has expected infinite flow, and the result follows by the preceding case. ■

C. Necessary and Sufficient Conditions for Ergodicity

Here, we provide the central result of this paper. The result establishes necessary and sufficient conditions for ergodicity of random models with weak feedback property and a common steady state $\pi > 0$ in expectation. The conditions are reliant on infinite flow, and guarantee that the

ergodicity of the model is equivalent to the ergodicity of the expected model. The result emerges as an outcome of several important results that we have developed so far. In particular, we combine the result $\mathcal{E} \subseteq \mathcal{F}$ stating that the ergodicity event is always contained in the infinite flow event (Theorem 1), the deterministic characterization of the infinite flow of Theorem 3, and the sufficient conditions of Theorem 6. We also make use of Theorem 4 providing conditions for equivalence of the ergodicity of the chain and the expected chain.

Theorem 7: (Infinite Flow Theorem) Let the random model $\{W(k)\}$ be independent, and have a common steady state $\pi > 0$ in expectation and weak feedback property. Then, the following conditions are equivalent:

- (a) The model is ergodic.
- (b) The model has infinite flow property.
- (c) The expected model has infinite flow property.
- (d) The expected model is ergodic.

Proof: First, we establish that parts (a), (b) and (c) are equivalent by showing that (a) \implies (b) \iff (c) \implies (a). In particular, by Theorem 1 we have $\mathcal{E} \subseteq \mathcal{F}$, showing that (a) \implies (b). By Theorem 3, parts (b) and (c) are equivalent. By Theorem 6, part (c) implies part (a). Now, we prove (a) \iff (d). Since (a) \iff (b), we have $\mathcal{E} = \mathcal{F}$ almost surely. Hence, by Theorem 4, the parts (a) and (d) are equivalent. ■

The infinite flow theorem combined with the deterministic characterization of the infinite flow model of Theorem 3 leads us to the following result.

Corollary 2: Let $\{A(k)\} \subset \mathbb{S}^m$ be a deterministic model that has a common steady state vector $\pi > 0$ and weak feedback property. Then, the chain $\{A(k)\}$ is ergodic if and only if $\sum_{k=0}^{\infty} A_S(k) = \infty$ for every nontrivial $S \subset [m]$.

Under the conditions of Theorem 7, the model admits consensus. This follows straightforwardly from relation $\mathcal{E} \subseteq \mathcal{C}$.

Corollary 3: Let the assumptions of Theorem 7 hold. Then, the model admits consensus.

The infinite flow theorem establishes the equivalence between the ergodicity of a chain and the expected chain for a class of independent random models. The crucial role in this result is played by the infinite flow and its equivalent deterministic characterization. Another crucial result is the interplay between the ergodicity and infinite flow of Theorem 4 yielding the equivalence between ergodicity of the chain and the expected chain. The following two examples are provided

to illustrate some straightforward applications of the infinite flow theorem.

I.I.D. Models. Consider an i.i.d. model $\{W(k)\}$. Then, the expected matrix $\bar{W} = \mathbb{E}[W(k)]$ is independent of k . Since \bar{W} is stochastic, we have $\pi^T \bar{W} = \pi^T$ for a stochastic vector $\pi \geq 0$. Therefore, i.i.d. model is an independent model with a common steady state π in expectation.

In [33], it is shown that for the class of i.i.d. models that have almost surely positive diagonal entries, the ergodicity of the expected model and the ergodicity of the original model are equivalent. The application of this result is reliant on the condition of the almost surely positive diagonal entries, which implies that the model has feedback property, as shown in Corollary 1. This property, however, is stronger than weak feedback property. At the same time, no requirement on the steady state vector is needed.

The application of the infinite flow theorem to the i.i.d. case would require weak feedback property and the existence of a steady state vector $\pi > 0$. Thus, there is a tradeoff in the conditions for the ergodicity provided by the infinite flow theorem and those given in [33]. To further illustrate the difference in the conditions, we consider the homogeneous deterministic model $\{A(k)\}$ of Example 5. The model $\{A(k)\}$ has weak feedback property and the steady state vector $\pi = \frac{1}{3}e$, so the ergodicity of the model can be deduced from the infinite flow theorem. At the same time, as seen in Example 5, the model does not have positive diagonal entries and, therefore, the ergodicity of the model cannot be deduced from the results in [33], [34]. In the light of this, the infinite flow theorem provides conditions for ergodicity that complement the conditions of [33], [34].

Gossip Algorithms on Time-varying Networks. As another application of the infinite flow theorem, we consider an extension of the standard gossip algorithm to time-varying networks. In particular, the gossip algorithm originally proposed in [5], [6] is for static networks. Here, we give a sufficient condition for the convergence of a gossip algorithm for networks with time-changing topology. Consider a network of m agents viewed as nodes of a graph with the node set $[m]$. Suppose that each agent has a private scalar value $x_i(0)$ at time $k = 0$. Now, let the interactions of the agents be random at nonnegative integer valued time instances k as follows: At any time $k \geq 1$, two different agents $i, j \in [m]$ wake up with probability $P_{ij}(k)$, where $P_{ij}(k) = P_{ji}(k)$ and $\sum_{i < j} P_{ij}(k) = 1$. Then, they set their values to the average of their current values, i.e., $x_i(k) = x_j(k) = \frac{1}{2}(x_i(k-1) + x_j(k-1))$. The choices of the pairs $\{i, j\}$ of interacting agents at different time instances are independent.

Based on the agent interaction model, define the independent random model $\{W(k)\}$ by:

$$W(k) = I - \frac{1}{2}(e_i - e_j)(e_i - e_j)^T \text{ with probability } P_{ij}(k). \quad (20)$$

Then, the dynamic system (1) driven by the random chain $\{W(k)\}$ describes the evolution of the vector $x(k)$ that has its i -th component value equal to agent i value, $x_i(k)$. As seen from (20), any realization of the model $\{W(k)\}$ is doubly stochastic. Hence, the model has a common steady state $\pi = \frac{1}{m}e$ in expectation. Also, the model has strong feedback property (with $\gamma = \frac{1}{2}$).

Lemma 13: For extended gossip algorithm (20), the consensus is almost sure if $\sum_{k=0}^{\infty} P_S(k) = \infty$ for any nontrivial set $S \subset [m]$.

Proof: The extended gossip algorithm satisfies the assumption of the infinite flow Theorem 7. Since $E[W_{ij}(k) + W_{ji}(k)] = P_{ij}(k)$ for all $i \neq j$ and $k \geq 0$, it follows that the model is ergodic if and only if $\sum_{k=0}^{\infty} P_S(k) = \sum_{k=0}^{\infty} \sum_{i \in S, j \in \bar{S}} P_{ij}(k) = \infty$. Thus, the model admits consensus if the preceding condition holds. ■

When $P(k) = P$ for all k as in [5], [6], we can consider the graph $G = ([m], E)$ where the edge $\{i, j\} \in E$ iff $P_{ij} = P_{ji} > 0$. In this case, it can be seen that the condition of Lemma 13 is equivalent to the requirement that the graph G is connected. One can further modify the algorithm in (20) to allow for time-varying weights, i.e., $x_i(k) = a(k)x_i(k-1) + (1-a(k))x_j(k-1)$ and $x_j(k) = a(k)x_j(k-1) + (1-a(k))x_i(k-1)$ with $a(k) \in (0, 1)$. Such a scheme is a natural generalization of the symmetric gossip model proposed in [13]. In this case, it can be verified that if $a(k) \in [a, 1-a]$ for $a \in (0, \frac{1}{2}]$, then the result of Lemma 13 still holds.

VI. CONCLUSION

We have studied the ergodicity and consensus problem for a linear discrete-time dynamic model driven by random stochastic matrices. We have introduced a concept of the infinite flow event and studied the relations among this event, ergodicity event and consensus event. The central result is the infinite flow theorem providing necessary and sufficient conditions for ergodicity of independent random models. The theorem captures the conditions ensuring the convergence of the random consensus algorithms, such as gossip and broadcast schemes [1], [11], [6]. Moreover, the infinite flow theorem captures simultaneously the conditions on the connectivity of the system and the sufficient information flow over time that have been important in studying the consensus and average consensus in deterministic settings [36], [37], [17], [28],

[24], [25], [26]. As illustrated briefly on two examples, the infinite flow theorem provides a convenient tool for studying the ergodicity of a model as well as consensus algorithms. Finally, we note that the work in this paper is readily extendible to the case when the initial state $x(0)$ in (1) is itself random and independent of the chain $\{W(k)\}$.

REFERENCES

- [1] T.C. Aysal, M.E. Yildriz, A.D. Sarwate, and A. Scaglione, *Broadcast gossip algorithms for consensus*, IEEE Transactions on Signal processing **57** (2009), 2748–2761.
- [2] D.P. Bertsekas and J.N. Tsitsiklis, *Parallel and distributed computation: Numerical methods*, Prentice-Hall Inc., 1989.
- [3] P. Billingsley, *Probability and measure*, John Wiley & Sons, Inc., USA, 1995.
- [4] P.A. Bliman, A. Nedić, and A. Ozdaglar, *Rate of convergence for consensus with delays*, Proceedings of the 47th IEEE Conference on Decision and Control, 2008.
- [5] S. Boyd, A. Ghosh, B. Prabhakar, and D. Shah, *Gossip algorithms: Design, analysis, and applications*, Proceedings IEEE Infocom, 2005, pp. 1653–1664.
- [6] ———, *Randomized gossip algorithms*, IEEE Transactions on Information Theory **52** (2006), no. 6, 2508–2530.
- [7] R. Carli, F. Fagnani, P. Frasca, T. Taylor, and S. Zampieri, *Average consensus on networks with transmission noise or quantization*, Proceedings of IEEE American Control Conference, 2007, pp. 4189–4194.
- [8] R. Carli, F. Fagnani, A. Speranzon, and S. Zampieri, *Communication constraints in the average consensus problem*, Automatica **44** (2008), no. 3, 671–684.
- [9] S. Chatterjee and E. Seneta, *Towards consensus: Some convergence theorems on repeated averaging*, Journal of Applied Probability **14** (1977), no. 1, 89–97.
- [10] R. Cogburn, *On products of random stochastic matrices*, In Random matrices and their applications (1986), 199–213.
- [11] A. Dimakis, A. Sarwate, and M. Wainwright, *Geographic gossip: Efficient averaging for sensor networks*, IEEE Transactions on Signal Processing **56** (2008), no. 3, 1205–1216.
- [12] R. Durrett, *Probability: Theory and examples*, third ed., Curt Hinrichs, 2005.
- [13] F. Fagnani and S. Zampieri, *Randomized consensus algorithms over large scale networks*, IEEE Journal on Selected Areas in Communications **26** (2008), no. 4, 634–649.
- [14] Y. Hatano, A.K. Das, and M. Mesbahi, *Agreement in presence of noise: Pseudogradients on random geometric networks*, Proceedings of the 44th IEEE Conference on Decision and Control, and European Control Conference, 2005, pp. 6382–6387.
- [15] M. Huang and J.H. Manton, *Stochastic approximation for consensus seeking: Mean square and almost sure convergence*, Proceedings of the 46th IEEE Conference on Decision and Control, 2007, pp. 306–311.
- [16] ———, *Coordination and consensus of networked agents with noisy measurements: Stochastic algorithms and asymptotic behavior*, SIAM Journal on Control and Optimization **48** (2009), no. 1, 134–161.
- [17] A. Jadbabaie, J. Lin, and S. Morse, *Coordination of groups of mobile autonomous agents using nearest neighbor rules*, IEEE Transactions on Automatic Control **48** (2003), no. 6, 988–1001.
- [18] S. Kar and J.M.F. Moura, *Distributed consensus algorithms in sensor networks: Link failures and channel noise*, IEEE Transactions on Signal Processing **57** (2009), no. 1, 355–369.
- [19] A. Kashyap, T. Basar, and R. Srikant, *Quantized consensus*, Automatica **43** (2007), no. 7, 1192–1203.

- [20] S. Li and T. Basar, *Asymptotic agreement and convergence of asynchronous stochastic algorithms*, IEEE Transactions on Automatic Control **32** (1987), no. 7, 612–618.
- [21] I. Lobel and A. Ozdaglar, *Distributed subgradient methods over random networks*, Tech. report, Laboratory for Information and Decision Systems, Report 2800, MIT, 2008.
- [22] K. Nawrotzki, *Discrete open systems on markov chains in a random environment. I*, Elektronische Informationsverarbeitung und Kybernetik **17** (1981), 569–599.
- [23] ———, *Discrete open systems on markov chains in a random environment. II*, Elektronische Informationsverarbeitung und Kybernetik **18** (1982), 83–98.
- [24] A. Nedić, A. Olshevsky, A. Ozdaglar, and J.N. Tsitsiklis, *On distributed averaging algorithms and quantization effects*, to appear in IEEE Transactions on Automatic Control, 2009.
- [25] A. Nedić and A. Ozdaglar, *On the rate of convergence of distributed subgradient methods for multi-agent optimization*, Proceedings of IEEE CDC, 2007, pp. 4711–4716.
- [26] ———, *Distributed subgradient methods for multi-agent optimization*, IEEE Transactions on Automatic Control **54** (2009), no. 1, 48–61.
- [27] R. Olfati-Saber and R.M. Murray, *Consensus problems in networks of agents with switching topology and time-delays*, IEEE Transactions on Automatic Control **49** (2004), no. 9, 1520–1533.
- [28] A. Olshevsky and J.N. Tsitsiklis, *Convergence speed in distributed consensus and averaging*, SIAM Journal on Control and Optimization **48** (2008), no. 1, 33–55.
- [29] S. Patterson, B. Bamieh, and A. El Abbadi, *Distributed average consensus with stochastic communication failures*, IEEE Transactions on Signal processing **57** (2009), 2748–2761.
- [30] P. Frasca, R. Carli, F. Fagnani and S. Zampieri, *Gossip consensus algorithms via quantized communication*, Automatica **46** (2010), no. 1, 70–80.
- [31] W. Ren and R.W. Beard, *Consensus seeking in multi-agent systems under dynamically changing interaction topologies*, IEEE Transactions on Automatic Control **50** (2005), no. 5, 655–661.
- [32] M. Rosenblatt, *Products of independent identically distributed stochastic matrices*, Journal of Journal of Mathematical Analysis and Applications **11** (1965), no. 1, 1–10.
- [33] A. Tahbaz-Salehi and A. Jadbabaie, *A necessary and sufficient condition for consensus over random networks*, IEEE Transactions on Automatic Control **53** (2008), no. 3, 791–795.
- [34] ———, *Consensus over ergodic stationary graph processes*, IEEE Transactions on Automatic Control **55** (2010), no. 1, 225–230.
- [35] B. Touri and A. Nedić, *Distributed consensus over network with noisy links*, Proceedings of the 12th International Conference on Information Fusion, 2009, pp. 146–154.
- [36] J.N. Tsitsiklis, *Problems in decentralized decision making and computation*, Ph.D. thesis, Dept. of Electrical Engineering and Computer Science, MIT, 1984.
- [37] J.N. Tsitsiklis and M. Athans, *Convergence and asymptotic agreement in distributed decision problems*, IEEE Transactions on Automatic Control **29** (1984), no. 1, 42–50.
- [38] J.N. Tsitsiklis, D.P. Bertsekas, and M. Athans, *Distributed asynchronous deterministic and stochastic gradient optimization algorithms*, IEEE Transactions on Automatic Control **31** (1986), no. 9, 803–812.